

³⁹Ar ATTA @ Heidelberg - the atom optical aspect

Florian Ritterbusch¹, Joachim Welte¹, Sven Ebser¹, Zhongy Feng¹, Werner Aeschbach-Hertig², Markus Oberthaler¹

¹ Kirchhoff-Institut für Physik, Heidelberg University, Heidelberg, Germany

² Institut für Umweltphysik, Heidelberg University, Heidelberg, Germany

The importance of ³⁹Ar as a dating tracer for the time range between 50 and 1000 years has clearly been identified [1]. The detection technique Atom Trap Trace Analysis (ATTA) has been realized for rare krypton isotopes in the past decade [2,3] and its principle applicability to ³⁹Ar has been demonstrated [4]. ATTA has the potential to measure ³⁹Ar on small samples down to less than 1 ccSTP of Argon, corresponding to about 100 ml of air, 2.5 l of water or 1 kg of ice. This would especially enable the application of ³⁹Ar as a tracer in oceanography or glaciology where samples smaller than 10 l of water or 1 kg of ice respectively are required for practical dating purposes.

We report on the development of an ATTA-setup for ³⁹Ar [5] which gives a ³⁹Ar-count rate of 0.6atoms/h for atmospheric samples. Initially, spectroscopy of the relevant hyperfine transition in ³⁹Ar has been performed [6,7]. Based on these results different atom-optical techniques have been developed to achieve the high ³⁹Ar-count rate. Especially, an additional deceleration stage behind the magneto-optical trap center was found to enhance the counting rate by a factor ~4. The ⁴⁰Ar-background could successfully be eliminated by selectively removing ⁴⁰Ar atoms with an additional laser frequency and thus allows the detection of single atoms with high signal to noise ratio without switching the trap parameters. A further increase in ³⁹Ar-count rate is intended by systematic studies on enriched samples including the hyperfine nature of ³⁹Ar, which is not present in the abundant argon isotopes.

[1] Loosli, H. H. (1983), A dating method with ³⁹Ar, *Earth and Planetary Science Letters*, 63, 51-62.

[2] C. Y. Chen, Y. M. Li, K. Bailey, T. P. O'Connor, L. Young, and Z.-T. Lu. *Ultrasensitive Isotope Trace Analyses with a Magneto-Optical Trap*. *Science*, 286(5442):1139-1141, 1999.

[3] N. C. Sturchio et al., One million year old groundwater in the Sahara revealed by krypton-81 and chlorine-36. *Geophys. Res. Lett.*, 31:L05503, 2004.

[4] Jiang, W. et al. (2011), ³⁹Ar detection at the 10⁻¹⁶ isotopic abundance level with atom trap trace analysis, *Phys. Rev. Lett.* 106, DOI: 10.1103/PhysRevLett.106.103001.

[5] J. Welte, F. Ritterbusch, I. Steinke, M. Henrich, W. Aeschbach-Hertig, and M. K. Oberthaler. *Toward the realization of atom trap trace analysis for ³⁹Ar*. *New Journal of Physics*, 2010.

[6] J. Welte, I. Steinke, M. Henrich, F. Ritterbusch, M. K. Oberthaler, W. Aeschbach-Hertig, W. H. Schwarz, and M. Trieloff. *Hyperfine spectroscopy of the 1s5 - 2p9 transition of ³⁹Ar*. *Review of Scientific Instruments*, 80(11):113109, 2009.

[7] W. Williams, Z.-T. Lu, K. Rudinger, C.-Y. Xu, R. Yokochi, and P. Müller. *Spectroscopic study of the cycling transition 4s[3/2]2-4p[5/2]3 at 811.8 nm in ³⁹Ar: Hyperfine structure and isotope shift*. *Phys. Rev. A*, 83:012512, 2011

[6] J. Welte, F. Ritterbusch, I. Steinke, M. Henrich, W. Aeschbach-Hertig, and M. K. Oberthaler. Toward the realization of atom trap trace analysis for ^{39}Ar . *New Journal of Physics*, 2010.